

Global temperature change

James Hansen^{*†}, Makiko Sato^{*†}, Reto Ruedy^{*§}, Ken Lo^{*§}, David W. Lea[¶], and Martin Medina-Elizade[¶]

^{*}National Aeronautics and Space Administration Goddard Institute for Space Studies, [†]Columbia University Earth Institute, and [§]Sigma Space Partners, Inc., 2880 Broadway, New York, NY 10025; and [¶]Department of Earth Science, University of California, Santa Barbara, CA 93106

Contributed by James Hansen, July 31, 2006

Global surface temperature has increased $\approx 0.2^\circ\text{C}$ per decade in the past 30 years, similar to the warming rate predicted in the 1980s in initial global climate model simulations with transient greenhouse gas changes. Warming is larger in the Western Equatorial Pacific than in the Eastern Equatorial Pacific over the past century, and we suggest that the increased West–East temperature gradient may have increased the likelihood of strong El Niños, such as those of 1983 and 1998. Comparison of measured sea surface temperatures in the Western Pacific with paleoclimate data suggests that this critical ocean region, and probably the planet as a whole, is approximately as warm now as at the Holocene maximum and within $\approx 1^\circ\text{C}$ of the maximum temperature of the past million years. We conclude that global warming of more than $\approx 1^\circ\text{C}$, relative to 2000, will constitute “dangerous” climate change as judged from likely effects on sea level and extermination of species.

climate change | El Niños | global warming | sea level | species extinctions

Global temperature is a popular metric for summarizing the state of global climate. Climate effects are felt locally, but the global distribution of climate response to many global climate forcings is reasonably congruent in climate models (1), suggesting that the global metric is surprisingly useful. We will argue further, consistent with earlier discussion (2, 3), that measurements in the Western Pacific and Indian Oceans provide a good indication of global temperature change.

We first update our analysis of surface temperature change based on instrumental data and compare observed temperature change with predictions of global climate change made in the 1980s. We then examine current temperature anomalies in the tropical Pacific Ocean and discuss their possible significance. Finally, we compare paleoclimate and recent data, using the Earth’s history to estimate the magnitude of global warming that is likely to constitute dangerous human-made climate change.

Modern Global Temperature Change

Global surface temperature in more than a century of instrumental data is recorded in the Goddard Institute for Space Studies analysis for 2005. Our analysis, summarized in Fig. 1, uses documented procedures for data over land (4), satellite measurements of sea surface temperature (SST) since 1982 (5), and a ship-based analysis for earlier years (6). Estimated 2σ error (95% confidence) in comparing nearby years of global temperature (Fig. 1A), such as 1998 and 2005, decreases from 0.1°C at the beginning of the 20th century to 0.05°C in recent decades (4). Error sources include incomplete station coverage, quantified by sampling a model-generated data set with realistic variability at actual station locations (7), and partly subjective estimates of data quality problems (8). The estimated uncertainty of global mean temperature implies that we can only state that 2005 was probably the warmest year.

The map of temperature anomalies for the first half-decade of the 21st century (Fig. 1B), relative to 1951–1980 climatology, shows that current warmth is nearly ubiquitous, generally larger over land than over ocean, and largest at high latitudes in the Northern Hemisphere. Our ranking of 2005 as the warmest year depends on the positive polar anomalies, especially the unusual Arctic warmth. In calculating the global mean, we give full weight to all regions based on area. Meteorological stations are sparse in the Arctic, but the estimated strong warm anomaly there in 2005 is consistent with

record low sea ice concentration and Arctic temperature anomalies inferred from infrared satellite data (9).

Our analysis includes estimated temperature anomalies up to 1,200 km from the nearest measurement station (7). Resulting spatial extrapolations and interpolations of temperature anomalies usually are meaningful for seasonal and longer time scales at middle and high latitudes, where the spatial scale of anomalies is set by Rossby waves (7). Thus, we believe that the unusual Arctic warmth of 2005 is real. Other characteristics of our analysis method are summarized in *Supporting Text*, which is published as supporting information on the PNAS web site.

Independent analysis by the National Climate Data Center (www.ncdc.noaa.gov/oa/climate/research/2005/ann/global.html), using a “teleconnection” approach to fill in data sparse regions, also finds 2005 to be the warmest year. The joint analysis of the University of East Anglia and the Hadley Centre (www.met-office.gov.uk/research/hadleycentre/obsdata/globaltemperature.html) also yields high global temperature for 2005, but a few hundredths of a degree cooler than in 1998.

Record, or near record, warmth in 2005 is notable, because global temperature did not receive a boost from an El Niño in 2005. The temperature in 1998, on the contrary, was lifted 0.2°C above the trend line by a “super El Niño” (see below), the strongest El Niño of the past century.

Global warming is now 0.6°C in the past three decades and 0.8°C in the past century. It is no longer correct to say “most global warming occurred before 1940.” A better summary is: slow global warming, with large fluctuations, over the century up to 1975, followed by rapid warming at a rate $\approx 0.2^\circ\text{C}$ per decade. Global warming was $\approx 0.7^\circ\text{C}$ between the late 19th century (the earliest time at which global mean temperature can be accurately defined) and 2000, and continued warming in the first half decade of the 21st century is consistent with the recent rate of $+0.2^\circ\text{C}$ per decade.

The conclusion that global warming is a real climate change, not an artifact due to measurements in urban areas, is confirmed by surface temperature change inferred from borehole temperature profiles at remote locations, the rate of retreat of alpine glaciers around the world, and progressively earlier breakup of ice on rivers and lakes (10). The geographical distribution of warming (Fig. 1B) provides further proof of real climate change. Largest warming is in remote regions including high latitudes. Warming occurs over ocean areas, far from direct human effects, with warming over ocean less than over land, an expected result for a forced climate change because of the ocean’s great thermal inertia.

Early Climate Change Predictions. Manabe and Wetherald (11) made the first global climate model (GCM) calculations of warming due

Author contributions: D.W.L. and M.M.-E. contributed data; J.H., M.S., R.R., K.L., D.W.L., and M.M.-E. analyzed data; and J.H. wrote the paper.

The authors declare no conflict of interest.


Freely available online through the PNAS open access option.

Abbreviations: SST, sea surface temperature; GHG, greenhouse gas; EEP, Eastern Equatorial Pacific; WEP, Western Equatorial Pacific; DAI, dangerous anthropogenic interference; BAU, business as usual; AS, alternative scenario; BC, black carbon.

[†]To whom correspondence should be addressed: E-mail: jhansen@giss.nasa.gov.

© 2006 by The National Academy of Sciences of the USA

A SST Change (°C) from 1870-1900 to 2001-2005



B Western and Eastern Pacific Temperature Anomalies (°C)




Fig. 3. Comparison of SST in West and East Equatorial Pacific Ocean. (A) SST in 2001–2005 relative to 1870–1900, from concatenation of two data sets (5, 6), as described in the text. (B) SSTs (12-month running means) in WEP and EEP relative to 1870–1900 means.

cover longer periods. Nevertheless, it is apparent that the first transient climate simulations (12) proved to be quite accurate, certainly not “wrong by 300%” (14). The assertion of 300% error may have been based on an earlier arbitrary comparison of 1988–1997 observed temperature change with only scenario A (18). Observed warming was slight in that 9-year period, which is too brief for meaningful comparison.

Super El Niños. The 1983 and 1998 El Niños were successively labeled “El Niño of the century,” because the warming in the Eastern Equatorial Pacific (EEP) was unprecedented in 100 years (Fig. 3). We suggest that warming of the Western Equatorial Pacific (WEP), and the absence of comparable warming in the EEP, has increased the likelihood of such “super El Niños.”

In the “normal” (La Niña) phase of El Niño Southern Oscillation the east-to-west trade winds push warm equatorial surface water to the west such that the warmest SSTs are located in the WEP near Indonesia. In this normal state, the thermocline is shallow in the EEP, where upwelling of cold deep water occurs, and deep in the WEP (figure 2 of ref. 20). Associated with this tropical SST gradient is a longitudinal circulation pattern in the atmosphere, the Walker cell, with rising air and heavy rainfall in the WEP and sinking air and drier conditions in the EEP. The Walker circulation enhances upwelling of cold water in the East Pacific, causing a powerful positive feedback, the Bjerknes (21) feedback, which tends to maintain the La Niña phase, as the SST gradient and resulting higher pressure in the EEP support east-to-west trade winds.

This normal state is occasionally upset when, by chance, the east-to-west trade winds slacken, allowing warm water piled up in the west to slosh back toward South America. If the fluctuation is large enough, the Walker circulation breaks down and the Bjerknes feedback loses power. As the east-to-west winds weaken, the Bjerknes feedback works in reverse, and warm waters move more strongly toward South America, reducing the thermocline tilt and cutting off upwelling of cold water along the South American coast. In this way, a classical El Niño is born.

Theory does not provide a clear answer about the effect of global warming on El Niños (19, 20). Most climate models yield either a tendency toward a more El Niño-like state or no clear change (22). It has been hypothesized that, during the early Pliocene, when the Earth was 3°C warmer than today, a permanent El Niño condition existed (23).

We suggest, on empirical grounds, that a near-term global warming effect is an increased likelihood of strong El Niños. Fig. 1B shows an absence of warming in recent years relative to 1951–1980 in the equatorial upwelling region off the coast of South America. This is also true relative to the earliest period of SST data, 1870–1900 (Fig. 3A). Fig. 7, which is published as supporting information on the PNAS web site, finds a similar result for linear

trends of SSTs. The trend of temperature minima in the East Pacific, more relevant for our purpose, also shows no equatorial warming in the East Pacific.

The absence of warming in the EEP suggests that upwelling water there is not yet affected much by global warming. Warming in the WEP, on the other hand, is 0.5–1°C (Fig. 3). We suggest that increased temperature difference between the near-equatorial WEP and EEP allows the possibility of increased temperature swing from a La Niña phase to El Niño, and that this is a consequence of global warming affecting the WEP surface sooner than it affects the deeper ocean.

Fig. 3B compares SST anomalies (12-month running means) in the WEP and EEP at sites (marked by circles in Fig. 3A) of paleoclimate data discussed below. Absolute temperatures at these sites are provided in Fig. 8, which is published as supporting information on the PNAS web site. Even though these sites do not have the largest warming in the WEP or largest cooling in the EEP, Fig. 3B reveals warming of the WEP relative to the EEP [135-year changes, based on linear trends, are +0.27°C (WEP) and –0.01°C (EEP)].

The 1983 and 1998 El Niños in Fig. 3B are notably stronger than earlier El Niños. This may be partly an artifact of sparse early data or the location of data sites, e.g., the late 1870s El Niño is relatively stronger if averages are taken over Niño 3 or a 5° × 10° box. Nevertheless, “super El Niños” clearly were more abundant in the last quarter of the 20th century than earlier in the century.

Global warming is expected to slow the mean tropical circulation (24–26), including the Walker cell. Sea level pressure data suggest a slowdown of the longitudinal wind by ≈3.5% in the past century (26). A relaxed longitudinal wind should reduce the WEP–EEP temperature difference on the broad latitudinal scale (≈10°N to 15°S) of the atmospheric Walker cell. Observed SST anomalies are consistent with this expectation, because the cooling in the EEP relative to WEP decreases at latitudes away from the narrower region strongly affected by upwelling off the coast of Peru (Fig. 3A). Averaged over 10°N to 15°S, observed warming is as great in the EEP as in the WEP (see also Fig. 7).

We make no suggestion about changes of El Niño frequency, and we note that an abnormally warm WEP does not assure a strong El Niño. The origin and nature of El Niños is affected by chaotic ocean and atmosphere variations, the season of the driving anomaly, the state of the thermocline, and other factors, assuring that there will always be great variability of strength among El Niños.

Will increased contrast between near-equatorial WEP and EEP SSTs be maintained or even increase with more global warming? The WEP should respond relatively rapidly to increasing GHGs. In the EEP, to the extent that upwelling water has not been exposed to the surface in recent decades, little warming is expected, and the

Criteria for Dangerous Warming. The United Nations Framework Convention on Climate Change (www.unfccc.int) has the objective “to achieve stabilization of GHG concentrations” at a level preventing “dangerous anthropogenic interference” (DAI) with climate, but climate change constituting DAI is undefined. We suggest that global temperature is a useful metric to assess proximity to DAI, because, with knowledge of the Earth’s history, global temperature can be related to principal dangers that the Earth faces.

We propose that two foci in defining DAI should be sea level and extinction of species, because of their potential tragic consequences and practical irreversibility on human time scales. In considering these topics, we find it useful to contrast two distinct scenarios abbreviated as “business-as-usual” (BAU) and the “alternative scenario” (AS).

BAU has growth of climate forcings as in intermediate or strong Intergovernmental Panel on Climate Change scenarios, such as A1B or A2 (10). CO₂ emissions in BAU scenarios continue to grow at ≈2% per year in the first half of this century, and non-CO₂ positive forcings such as CH₄, N₂O, O₃, and black carbon (BC) aerosols also continue to grow (10). BAU, with nominal climate sensitivity ($3 \pm 1^\circ\text{C}$ for doubled CO₂), yields global warming (above year 2000 level) of at least 2–3°C by 2100 (10, 17).

AS has declining CO₂ emissions and an absolute decrease of non-CO₂ climate forcings, chosen such that, with nominal climate sensitivity, global warming (above year 2000) remains <1°C. For example, one specific combination of forcings has CO₂ peaking at 475 ppm in 2100 and a decrease of CH₄, O₃, and BC sufficient to balance positive forcing from increase of N₂O and decrease of sulfate aerosols. If CH₄, O₃, and BC do not decrease, the CO₂ cap in AS must be lower.

Sea level implications of BAU and AS scenarios can be considered in two parts: equilibrium (long-term) sea level change and ice sheet response time. Global warming <1°C in AS keeps temperatures near the peak of the warmest interglacial periods of the past million years. Sea level may have been a few meters higher than today in some of those periods (10). In contrast, sea level was 25–35 m higher the last time that the Earth was 2–3°C warmer than today, i.e., during the Middle Pliocene about three million years ago (32).

Ice sheet response time can be investigated from paleoclimate evidence, but inferences are limited by imprecise dating of climate and sea level changes and by the slow pace of weak paleoclimate forcings compared with stronger rapidly increasing human-made forcings. Sea level rise lagged tropical temperature by a few thousand years in some cases (28), but in others, such as Meltwater Pulse 1A ≈14,000 years ago (33), sea level rise and tropical temperature increase were nearly synchronous. Intergovernmental Panel on Climate Change (10) assumes negligible contribution to 2100 sea level change from loss of Greenland and Antarctic ice, but that conclusion is implausible (17, 34). BAU warming of 2–3°C would bathe most of Greenland and West Antarctic in melt-water during lengthened melt seasons. Multiple positive feedbacks, including reduced surface albedo, loss of buttressing ice shelves, dynamical response of ice streams to increased melt-water, and lowered ice surface altitude would assure a large fraction of the equilibrium ice sheet response within a few centuries, at most (34).

Sea level rise could be substantial even in the AS, ≈1 m per century, and cause problems for humanity due to high population in coastal areas (10). However, AS problems would be dwarfed by the disastrous BAU, which could yield sea level rise of several meters per century with eventual rise of tens of meters, enough to transform global coastlines.

Extinction of animal and plant species presents a picture analogous to that for sea level. Extinctions are already occurring as a result of various stresses, mostly human-made, including climate change (35). Plant and animal distributions are a reflection of the regional climates to which they are adapted. Thus, plants and animals attempt to migrate in response to climate change, but their

Poleward Migration Rate of Isotherms (km/decade)




Fig. 6. Poleward migration rate of isotherms in surface observations (A and B) and in climate model simulations (17) for 2000–2100 for Intergovernmental Panel on Climate Change scenario A2 (10) and an alternative scenario of forcings that keeps global warming after 2000 less than 1°C (17) (C and D). Numbers in upper right are global means excluding the tropical band.

paths may be blocked by human-constructed obstacles or natural barriers such as coastlines.

A study of 1,700 biological species (36) found poleward migration of 6 km per decade and vertical migration in alpine regions of 6 m per decade in the second half of the 20th century, within a factor of two of the average poleward migration rate of surface isotherms (Fig. 6A) during 1950–1995. More rapid warming in 1975–2005 yields an average isotherm migration rate of 40 km per decade in the Northern Hemisphere (Fig. 6B), exceeding known paleoclimate rates of change. Some species are less mobile than others, and ecosystems involve interactions among species, so such rates of climate change, along with habitat loss and fragmentation, new invasive species, and other stresses are expected to have severe impact on species survival (37).

The total distance of isotherm migration, as well as migration rate, affects species survival. Extinction is likely if the migration distance exceeds the size of the natural habitat or remaining habitat fragment. Fig. 6 shows that the 21st century migration distance for a BAU scenario (≈600 km) greatly exceeds the average migration distance for the AS (≈100 km).

It has been estimated (38) that a BAU global warming of 3°C over the 21st century could eliminate a majority (≈60%) of species on the planet. That projection is not inconsistent with mid-century BAU effects in another study (37) or scenario sensitivity of stress effects (35). Moreover, in the Earth’s history several mass extinctions of 50–90% of species have accompanied global temperature changes of ≈5°C (39).

We infer that even AS climate change, which would slow warming to <0.1°C per decade over the century, would contribute to species loss that is already occurring due to a variety of stresses. However, species loss under BAU has the potential to be truly disastrous, conceivably with a majority of today’s plants and animals headed toward extermination.

Discussion

The pattern of global warming (Fig. 1B) has assumed expected characteristics, with high latitude amplification and larger warming over land than over ocean, as GHGs have become the dominant climate forcing in recent decades. This pattern results mainly from the ice–snow albedo feedback and the response times of ocean and land.

